arXiv:1107.2488v1 [cond-mat.mtrl-sci] 13 Jul 2011

The relationship between the density functional Hartree-plus-exchange-correlation potential for an integer N-electron and $(N-1)$ electron system.

Daniel P. Joubert[1, a)]

Centre for Theoretical Physics, University of the Witwatersrand, PO Wits 2050, Johannesburg, South Africa

(Dated: 14 July 2011)

It is shown that the Hartree-plus-exchange-correlation density functional potential for an integer N-electron system differs by a constant form the corresponding potential for an $(N-1)$-electron system if the densities are determined from the same external potential.

PACS numbers: 71.15.Mb, 31.15.E , 71.45.Gm
Keywords: density functional, exchange-correlation

---

a)Electronic mail: daniel.joubert2@wits.ac.za

I. INTRODUCTION

In the early 1980's it was discovered that the density functional exchange correlation potential has a derivative discontinuity when the particle number crosses and integer[1,2]. In this paper a relationship between the potentials of systems with different particle numbers is proven. It will be shown that the Hartree-plus-exchange-correlation potential for an integer N-electron ($N > 2$) system differs by a constant from the corresponding potential of an $(N-1)$-electron system if the densities are determined from the same external potential. As a corollary it follows that the functional derivative of the independent particle kinetic energy functional of the N and $(N-1)$-electron systems also differ by a constant.

II. PROOF

In the adiabatic connection approach[3–6] of the constrained minimization formulation of density functional theory[7–10] the Hamiltonian $\hat{H}^{\gamma}$ for a system of N electrons is given by

$$\hat{H}_N^{\gamma} = \hat{T}^N + \gamma \hat{V}_{\text{ee}}^N + \hat{v}_{N,\text{ext}}^{\gamma}\left[\rho_N\right]. \tag{1}$$

Atomic units, $\hbar = e = m = 1$ are used throughout. $\hat{T}$ is the kinetic energy operator,

$$\hat{T}^N = -\frac{1}{2}\sum_{i=1}^{N} \nabla_i^2, \tag{2}$$

and $\gamma \hat{V}_{\text{ee}}$ is a scaled electron-electron interaction,

$$\gamma \hat{V}_{\text{ee}}^N = \gamma \sum_{i<j}^{N} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}. \tag{3}$$

The the external potential

$$\hat{v}_{N,\text{ext}}^{\gamma}\left[\rho_N\right] = \sum_{i=1}^{N} v_{\text{ext}}^{\gamma}\left(\left[\rho_N\right];\mathbf{r}_i\right), \tag{4}$$

is constructed to keep the charge density fixed at $\rho_N(\mathbf{r})$,the ground state charge density of the fully interacting system ($\gamma = 1$), for all values of the coupling constant γ. It has the form[10,11]

$$\begin{aligned} v_{\text{ext}}^{\gamma}(\left[\rho_N\right];\mathbf{r}) = {} & (1-\gamma)\, v_{ux}([\rho_N];\mathbf{r}) \\ & + v_c^1([\rho_N];\mathbf{r}) - v_c^{\gamma}([\rho_N];\mathbf{r}) + v_{\text{ext}}^1(\left[\rho_N\right];\mathbf{r}), \end{aligned} \tag{5}$$

where $v^1_{\text{ext}}([\rho_N]\,;\mathbf{r}) = v_{\text{ext}}(\mathbf{r})$ is the external potential at full coupling strength, $\gamma = 1$, and $v^0_{\text{ext}}([\rho_N]\,;\mathbf{r})$ is non-interacting Kohn-Sham potential. The exchange plus Hartree potential[12,13]$v_{ux}([\rho_N];\mathbf{r})$, is independent of γ, while the correlation potential $v^{\gamma}_c([\rho_N];\mathbf{r})$ depends in the scaling parameter γ.

The chemical potential

$$\mu = E^{\gamma}_N - E^{\gamma}_{N-1} \tag{6}$$

depends on the asymptotic decay of the charge density[12–14], and hence is independent of the coupling constant γ[15,16]. In Eq. (6) E^{γ}_{N-1} is the groundstate energy of the $(N-1)$-electron system with the same single-particle external potential $v^{\gamma}_{\text{ext}}([\rho_N]\,;\mathbf{r})$ as the N-electron system:

$$\begin{aligned}
\hat{H}^{\gamma}_{N-1}\left|\Psi^{\gamma}_{\rho^{\gamma}_{N-1}}\right\rangle &= E^{\gamma}_{N-1}\left|\Psi^{\gamma}_{\rho^{\gamma}_{N-1}}\right\rangle \\
\hat{H}^{\gamma}_{N-1} &= \hat{T}^{N-1} + \gamma\hat{V}^{N-1}_{\text{ee}} + \hat{v}^{\gamma}_{N-1,\text{ext}}[\rho_N] \\
\hat{v}^{\gamma}_{N-1,\text{ext}}[\rho_N] &= \sum_{i=1}^{N-1} v^{\gamma}_{\text{ext}}([\rho_N]\,;\mathbf{r}_i)
\end{aligned} \tag{7}$$

Note that by construction of $v^{\gamma}_{\text{ext}}([\rho_N]\,;\mathbf{r})$, Eq. (5) ρ_N is independent of γ, but the groundstate density of the $(N-1)$-electron system ρ^{γ}_{N-1}, is a function of γ.

The correlation energy $E^{\gamma}_c\left[\rho^{\gamma}_{N-1}\right]$ is defined as[10]

$$\begin{aligned}
E^{\gamma}_c\left[\rho^{\gamma}_{N-1}\right] = &\left\langle\Psi^{\gamma}_{\rho^{\gamma}_{N-1}}\left|\hat{T}^{N-1} + \gamma\hat{V}^{N-1}_{\text{ee}}\right|\Psi^{\gamma}_{\rho^{\gamma}_{N-1}}\right\rangle \\
&- \left\langle\Psi^{0}_{\rho^{\gamma}_{N-1}}\left|\hat{T}^{N-1} + \gamma\hat{V}^{N-1}_{\text{ee}}\right|\Psi^{0}_{\rho^{\gamma}_{N-1}}\right\rangle,
\end{aligned} \tag{8}$$

where $\left|\Psi^{0}_{\rho^{\gamma}_{N-1}}\right\rangle$ is the Kohn-Sham $(N-1)$ independent particle groundstate wavefunction that yields the same density as the interacting $(N-1)$-electron system at coupling strength γ. Since the correlation part of the kinetic energy is given by

$$\begin{aligned}
&T^{\gamma}_c\left[\rho^{\gamma}_{N-1}\right] \\
&= \left\langle\Psi^{\gamma}_{\rho^{\gamma}_{N-1}}\left|\hat{T}^{N-1}\right|\Psi^{\gamma}_{\rho^{\gamma}_{N-1}}\right\rangle - \left\langle\Psi^{0}_{\rho^{\gamma}_{N-1}}\left|\hat{T}^{N-1}\right|\Psi^{0}_{\rho^{\gamma}_{N-1}}\right\rangle,
\end{aligned} \tag{9}$$

and hence

$$\begin{aligned}
&\frac{E^{\gamma}_c\left[\rho^{\gamma}_{N-1}\right] - T^{\gamma}_c\left[\rho^{\gamma}_{N-1}\right]}{\gamma} \\
&= \left\langle\Psi^{\gamma}_{\rho^{\gamma}_{N-1}}\left|\hat{V}^{N-1}_{\text{ee}}\right|\Psi^{\gamma}_{\rho^{\gamma}_{N-1}}\right\rangle - \left\langle\Psi^{0}_{\rho^{\gamma}_{N-1}}\left|\hat{V}^{N-1}_{\text{ee}}\right|\Psi^{0}_{\rho^{\gamma}_{N-1}}\right\rangle,
\end{aligned} \tag{10}$$

the derivative of $E_c^\gamma\left[\rho_{N-1}^\gamma\right]$ with respect to γ can be expressed as

$$\begin{aligned}
&\frac{\partial}{\partial\gamma}E_c^\gamma\left[\rho_{N-1}^\gamma\right] \\
&= \frac{E_c^\gamma\left[\rho_{N-1}^\gamma\right] - T_c^\gamma\left[\rho_{N-1}^\gamma\right]}{\gamma} \\
&\quad + \left\langle \frac{\partial}{\partial\gamma}\Psi^\gamma_{\rho^\gamma_{N-1}} \left| \hat{T}^{N-1} + \gamma\hat{V}_{\text{ee}}^{N-1} \right| \Psi^\gamma_{\rho^\gamma_{N-1}} \right\rangle \\
&\quad + \left\langle \Psi^\gamma_{\rho^\gamma_{N-1}} \left| \hat{T}^{N-1} + \gamma\hat{V}_{\text{ee}}^{N-1} \right| \frac{\partial}{\partial\gamma}\Psi^\gamma_{\rho^\gamma_{N-1}} \right\rangle \\
&\quad - \left\langle \frac{\partial}{\partial\gamma}\Psi^0_{\rho^\gamma_{N-1}} \left| \hat{T}^{N-1} + \gamma\hat{V}_{\text{ee}}^{N-1} \right| \Psi^0_{\rho^\gamma_{N-1}} \right\rangle \\
&\quad - \left\langle \Psi^0_{\rho^\gamma_{N-1}} \left| \hat{T}^{N-1} + \gamma\hat{V}_{\text{ee}}^{N-1} \right| \frac{\partial}{\partial\gamma}\Psi^0_{\rho^\gamma_{N-1}} \right\rangle .
\end{aligned} \tag{11}$$

Upon adding and subtracting (c.c. stands for the complex conjugate of the previous term)

$$\begin{gathered}
\left(\left\langle \frac{\partial}{\partial\gamma}\Psi^\gamma_{\rho^\gamma_{N-1}} \left| \hat{v}^\gamma_{N-1,\text{ext}}\left[\rho_N\right] \right| \Psi^\gamma_{\rho^\gamma_{N-1}} \right\rangle + \text{c.c}\right) + \\
\left(\left\langle \frac{\partial}{\partial\gamma}\Psi^\gamma_{\rho^\gamma_{N-1}} \right|_{\gamma=0} \left| \hat{v}^0_{N-1,\text{ext}}\left[\rho_N\right] \right| \Psi^0_{\rho^0_{N-1}} \right\rangle + \text{c.c}\right)
\end{gathered} \tag{12}$$

and utilizing the normalization of the wavefunctions which implies that

$$\frac{\partial}{\partial\gamma}\left\langle \Psi^\gamma_{\rho^\gamma_{N-1}} | \Psi^\gamma_{\rho^\gamma_{N-1}} \right\rangle = 0, \tag{13}$$

Eq. (11) becomes

$$\begin{aligned}
&\frac{\partial}{\partial\gamma}E_c^\gamma\left[\rho_{N-1}^\gamma\right] \\
&= \frac{E_c^\gamma\left[\rho_{N-1}^\gamma\right] - T_c^\gamma\left[\rho_{N-1}^\gamma\right]}{\gamma} \\
&\quad - \left\langle \frac{\partial}{\partial\gamma}\Psi^\gamma_{\rho^\gamma_{N-1}} \left| \hat{v}^\gamma_{N-1,\text{ext}}\left[\rho_N\right] \right| \Psi^\gamma_{\rho^\gamma_{N-1}} \right\rangle - \left\langle \Psi^\gamma_{\rho^\gamma_{N-1}} \left| \hat{v}^\gamma_{N-1,\text{ext}}\left[\rho_N\right] \right| \frac{\partial}{\partial\gamma}\Psi^\gamma_{\rho^\gamma_{N-1}} \right\rangle \\
&\quad + \left\langle \frac{\partial}{\partial\gamma}\Psi^0_{\rho^\gamma_{N-1}} \left| \hat{v}^0_{N-1,\text{ext}}\left[\rho_N\right] \right| \Psi^0_{\rho^\gamma_{N-1}} \right\rangle + \left\langle \Psi^0_{\rho^\gamma_{N-1}} \left| \hat{v}^0_{N-1,\text{ext}}\left[\rho_N\right] \right| \frac{\partial}{\partial\gamma}\Psi^0_{\rho^\gamma_{N-1}} \right\rangle \\
&\quad -\gamma\frac{\partial}{\partial\gamma}\left\langle \Psi^0_{\rho^\gamma_{N-1}} \left| \hat{V}_{\text{ee}}^{N-1} \right| \Psi^0_{\rho^\gamma_{N-1}} \right\rangle .
\end{aligned} \tag{14}$$

Now[12,13]

$$\begin{aligned}
\left\langle \Psi^0_{\rho^\gamma_{N-1}} \left| \hat{V}_{\text{ee}}^{N-1} \right| \Psi^0_{\rho^\gamma_{N-1}} \right\rangle &= E_x\left[\rho_{N-1}^\gamma\right] + U\left[\rho_{N-1}^\gamma\right] \\
&= E_{ux}\left[\rho_{N-1}^\gamma\right] .
\end{aligned} \tag{15}$$

is the sum of the exchange $E_x\left[\rho_{N-1}^{\gamma}\right]$ and mutual Coulomb interaction energy $U\left[\rho_{N-1}^{\gamma}\right]$ of the $(N-1)$-electron system. The charge density ρ_{N-1}^{γ} is a function of γ, therefore[12]

$$\frac{\partial}{\partial\gamma}\left\langle\Psi_{\rho_{N-1}^{\gamma}}^{0}\left|\hat{V}_{\text{ee}}^{N-1}\right|\Psi_{\rho_{N-1}^{\gamma}}^{0}\right\rangle=\int d^3r\frac{\partial\rho_{N-1}^{\gamma}(\mathbf{r})}{\partial\gamma}v_{ux}\left(\left[\rho_{N-1}^{\gamma}\right];\mathbf{r}\right), \tag{16}$$

where

$$v_{ux}\left(\left[\rho_{N-1}^{\gamma}\right];\mathbf{r}\right)=\frac{\delta}{\delta\rho_{N-1}^{\gamma}(\mathbf{r})}\left(E_x\left[\rho_{N-1}^{\gamma}\right]+U\left[\rho_{N-1}^{\gamma}\right]\right) \tag{17}$$

is the sum of the exchange and Hartree potentials for the $(N-1)$-electron system. Using Eq. (16), Eq. (5) and the fact that $\left|\Psi_{\rho_{N-1}^{\gamma}}^{\gamma}\right\rangle$ and $\left|\Psi_{\rho_{N-1}^{\gamma}}^{0}\right\rangle$ yield the same density ρ_{N-1}^{γ}, Eq. (14) can be cast as

$$\begin{aligned}&\frac{\partial}{\partial\gamma}E_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right]\\=&\frac{E_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right]-T_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right]}{\gamma}\\&+\int d^3r\frac{\partial\rho_{N-1}^{\gamma}(\mathbf{r})}{\partial\gamma}\left(v_c^{\gamma}\left(\left[\rho_N\right];\mathbf{r}\right)+\gamma v_{ux}\left(\left[\rho_N\right];\mathbf{r}\right)-\gamma v_{ux}\left(\left[\rho_{N-1}^{\gamma}\right];\mathbf{r}\right)\right)\end{aligned} \tag{18}$$

From the definition of $E_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right]$, Eq. (8), the correlation energy can also be written as

$$E_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right]=\hat{T}^{\gamma}\left[\rho_{N-1}^{\gamma}\right]-\hat{T}^{0}\left[\rho_{N-1}^{\gamma}\right]+\gamma\hat{V}_{\text{ee}}\left[\rho_{N-1}^{\gamma}\right]-\gamma E_{ux}\left[\rho_{N-1}^{\gamma}\right] \tag{19}$$

where

$$\hat{T}^{\gamma}\left[\rho_{N-1}^{\gamma}\right]=\left\langle\Psi_{\rho_{N-1}^{\gamma}}^{\gamma}\left|\hat{T}^{N-1}\right|\Psi_{\rho_{N-1}^{\gamma}}^{\gamma}\right\rangle$$

$$\hat{T}^{0}\left[\rho_{N-1}^{\gamma}\right]=\left\langle\Psi_{\rho_{N-1}^{\gamma}}^{0}\left|\hat{T}^{N-1}\right|\Psi_{\rho_{N-1}^{\gamma}}^{0}\right\rangle \tag{20}$$

$$.\hat{V}_{\text{ee}}\left[\rho_{N-1}^{\gamma}\right]=\left\langle\Psi_{\rho_{N-1}^{\gamma}}^{\gamma}\left|\hat{V}_{\text{ee}}^{N-1}\right|\Psi_{\rho_{N-1}^{\gamma}}^{\gamma}\right\rangle. \tag{21}$$

It now follows that

$$\begin{aligned}&\frac{\partial}{\partial\gamma}E_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right]\\=&\frac{E_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right]-T_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right]}{\gamma}\\&+\int d^3r\frac{\partial\rho_{N-1}^{\gamma}(\mathbf{r})}{\partial\gamma}\left(\frac{\delta T_{N-1}^{\gamma}\left[\rho_{N-1}^{\gamma}\right]}{\delta\rho_{N-1}^{\gamma}(\mathbf{r})}-\frac{\delta T_{N-1}^{0}\left[\rho_{N-1}^{\gamma}\right]}{\delta\rho_{N-1}^{\gamma}(\mathbf{r})}-\gamma\frac{\delta E_{ux}\left[\rho_{N-1}^{\gamma}\right]}{\delta\rho_{N-1}^{\gamma}(\mathbf{r})}\right)\\&+\gamma\frac{\partial}{\partial\gamma}V_{\text{ee}}\left[\rho_{N-1}^{\gamma}\right]\end{aligned} \tag{22}$$

where use was made of the relation

$$\frac{E_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right] - T_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right]}{\gamma} = \hat{V}_{\mathrm{ee}}\left[\rho_{N-1}^{\gamma}\right] - E_{ux}\left[\rho_{N-1}^{\gamma}\right]. \tag{23}$$

The last term in Eq. (22) can be transformed as follows:

$$\begin{aligned}
&\frac{\partial}{\partial\gamma} V_{\mathrm{ee}}\left[\rho_{N-1}^{\gamma}\right] \\
&= \left\langle \frac{\partial}{\partial\gamma}\Psi^{\gamma}_{\rho^{\gamma}_{N-1}} \middle| \hat{V}_{\mathrm{ee}}^{N-1} \middle| \Psi^{\gamma}_{\rho^{\gamma}_{N-1}} \right\rangle + \left\langle \Psi^{\gamma}_{\rho^{\gamma}_{N-1}} \middle| \hat{V}_{\mathrm{ee}}^{N-1} \middle| \frac{\partial}{\partial\gamma}\Psi^{\gamma}_{\rho^{\gamma}_{N-1}} \right\rangle \\
&= \frac{1}{\gamma}\left(\left\langle \frac{\partial}{\partial\gamma}\Psi^{\gamma}_{\rho^{\gamma}_{N-1}} \middle| \hat{T}^{N-1} + \gamma\hat{V}_{ee}^{N-1} + \hat{v}^{\gamma}_{N-1,\mathrm{ext}}\left[\rho_N\right] \middle| \Psi^{\gamma}_{\rho^{\gamma}_{N-1}} \right\rangle \right. \\
&\quad \left. + \left\langle \Psi^{\gamma}_{\rho^{\gamma}_{N-1}} \middle| \hat{T}^{N-1} + \gamma\hat{V}_{ee}^{N-1} + \hat{v}^{\gamma}_{N-1,\mathrm{ext}}\left[\rho_N\right] \middle| \frac{\partial}{\partial\gamma}\Psi^{\gamma}_{\rho^{\gamma}_{N-1}} \right\rangle \right) \\
&\quad -\frac{1}{\gamma}\left(\left\langle \frac{\partial}{\partial\gamma}\Psi^{\gamma}_{\rho^{\gamma}_{N-1}} \middle| \hat{T}^{N-1} + \hat{v}^{\gamma}_{N-1,\mathrm{ext}}\left[\rho_N\right] \middle| \Psi^{\gamma}_{\rho^{\gamma}_{N-1}} \right\rangle \right. \\
&\quad \left. + \left\langle \Psi^{\gamma}_{\rho^{\gamma}_{N-1}} \middle| \hat{T}^{N-1} + \hat{v}^{\gamma}_{N-1,\mathrm{ext}}\left[\rho_N\right] \middle| \frac{\partial}{\partial\gamma}\Psi^{\gamma}_{\rho^{\gamma}_{N-1}} \right\rangle \right) \\
&= -\frac{1}{\gamma}\int d^3r \frac{\partial\rho^{\gamma}_{N-1}(\mathbf{r})}{\partial\gamma}\left(\frac{\delta T^{\gamma}_{N-1}\left[\rho^{\gamma}_{N-1}\right]}{\delta\rho^{\gamma}_{N-1}(\mathbf{r})} + v^{\gamma}_{\mathrm{ext}}(\left[\rho_N\right];\mathbf{r}) \right).
\end{aligned} \tag{24}$$

Here use was made Eqs. (7) and (13). Combining Eqs. (22) and (24) leads to

$$\begin{aligned}
&\frac{\partial}{\partial\gamma} E_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right] \\
&= \frac{E_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right] - T_c^{\gamma}\left[\rho_{N-1}^{\gamma}\right]}{\gamma} \\
&\quad + \int d^3r \frac{\partial\rho^{\gamma}_{N-1}(\mathbf{r})}{\partial\gamma}\left(-v^{\gamma}_{\mathrm{ext}}(\left[\rho_N\right];\mathbf{r}) - \frac{\delta T^{0}_{N-1}\left[\rho^{\gamma}_{N-1}\right]}{\delta\rho^{\gamma}_{N-1}(\mathbf{r})} - \gamma v_{ux}\left(\left[\rho^{\gamma}_{N-1}\right];\mathbf{r}\right) \right).
\end{aligned} \tag{25}$$

At the solution point[12,13] the following equation has to be satisfied for the $(N-1)$-electron system:

$$\frac{\delta T^{0}_{N-1}\left[\rho^{\gamma}_{N-1}\right]}{\delta\rho^{\gamma}_{N-1}(\mathbf{r})} + \gamma v_{ux}\left(\left[\rho^{\gamma}_{N-1}\right];\mathbf{r}\right) + v_c^{\gamma}\left(\left[\rho^{\gamma}_{N-1}\right];\mathbf{r}\right) + v^{\gamma}_{\mathrm{ext}}(\left[\rho_N\right];\mathbf{r}) = \mu^{\gamma}_{N-1}, \tag{26}$$

where μ^{γ}_{N-1} is a constant, the chemical potential of the $(N-1)$-electrons moving in the external potential $v^{\gamma}_{\mathrm{ext}}(\left[\rho_N\right];\mathbf{r})$. Taking into account that the number of electrons is fixed at $(N-1)$ independent of γ, it follows that

$$\begin{aligned}
\int d^3r \frac{\partial\rho^{\gamma}_{N-1}(\mathbf{r})}{\partial\gamma} &= \frac{\partial}{\partial\gamma}(N-1) \\
&= 0,
\end{aligned} \tag{27}$$

where it is assumed that the order of the integration and derivative can be reversed. From Eqs. (25), (26) and (27)

$$\begin{aligned}&\frac{\partial}{\partial\gamma}E_c^\gamma\left[\rho_{N-1}^\gamma\right]\\&=\frac{E_c^\gamma\left[\rho_{N-1}^\gamma\right]-T_c^\gamma\left[\rho_{N-1}^\gamma\right]}{\gamma}+\int d^3r\frac{\partial\rho_{N-1}^\gamma\left(\mathbf{r}\right)}{\partial\gamma}v_c^\gamma\left(\left[\rho_{N-1}^\gamma\right];\mathbf{r}\right).\end{aligned} \tag{28}$$

Comparing Eqs. (18) and (28) shows that

$$0=\int d^3r\frac{\partial\rho_{N-1}^\gamma\left(\mathbf{r}\right)}{\partial\gamma}\left(v_{uxc}^\gamma\left(\left[\rho_N\right];\mathbf{r}\right)-v_{uxc}^\gamma\left(\left[\rho_{N-1}^\gamma\right];\mathbf{r}\right)\right) \tag{29}$$

where

$$v_{uxc}^\gamma\left(\left[\rho\right];\mathbf{r}\right)=v_c^\gamma\left(\left[\rho\right];\mathbf{r}\right)+\gamma v_{ux}\left(\left[\rho\right];\mathbf{r}\right) \tag{30}$$

is the Hartree plus exchange-correlation potential for a system with density ρ and coupling strength γ.

The charge density $\rho_{N-1}^\gamma\left(\mathbf{r}\right)$ is a functional of the potential $v_{\text{ext}}^\gamma\left(\left[\rho_N\right];\mathbf{r}\right)$ as can be seen from Eq. (7)[12,13]. Therefore

$$\begin{aligned}\frac{\partial\rho_{N-1}^\gamma\left(\mathbf{r}\right)}{\partial\gamma}&=\int d^3r'\left.\frac{\delta\rho_{N-1}^\gamma\left(\mathbf{r}\right)}{\delta v_{\text{ext}}^\gamma\left(\left[\rho_N\right];\mathbf{r}'\right)}\right|_{N-1}\frac{\partial}{\partial\gamma}v_{\text{ext}}^\gamma\left(\left[\rho_N\right];\mathbf{r}'\right)\\&=-\int d^3r'\left.\frac{\delta\rho_{N-1}^\gamma\left(\mathbf{r}\right)}{\delta v_{\text{ext}}^\gamma\left(\left[\rho_N\right];\mathbf{r}'\right)}\right|_{N-1}\left(v_{ux}([\rho_N];\mathbf{r}')+\frac{\partial}{\partial\gamma}v_c^\gamma([\rho_N];\mathbf{r}')\right).\end{aligned} \tag{31}$$

Now

$$\left.\frac{\delta\rho_{N-1}^\gamma\left(\mathbf{r}\right)}{\delta v_{\text{ext}}^\gamma\left(\left[\rho_N\right];\mathbf{r}'\right)}\right|_{N-1}=\chi_{N-1}^\gamma\left(\mathbf{r},\mathbf{r}'\right) \tag{32}$$

is the density response function of the $(N-1)$-particle system. From stability considerations $\chi_{N-1}^\gamma\left(\mathbf{r},\mathbf{r}'\right)$ is negative semi-definite and has one zero eigenvalue which corresponds to the invariance of the density when the potential is changed by a constant[12,13]. This implies that

$$\begin{aligned}0&=\int d^3r\frac{\partial\rho_{N-1}^\gamma\left(\mathbf{r}\right)}{\partial\gamma}f\left(\mathbf{r}\right)\\&=-\int d^3r\int d^3r'f\left(\mathbf{r}\right)\chi_{N-1}^\gamma\left(\mathbf{r},\mathbf{r}'\right)\left(v_{ux}([\rho_N];\mathbf{r}')+\frac{\partial}{\partial\gamma}v_c^\gamma([\rho_N];\mathbf{r}')\right)\end{aligned} \tag{33}$$

is only possible if $f\left(\mathbf{r}\right)=\text{constant}$ since $v_{ux}([\rho_N];\mathbf{r}')+\frac{\partial}{\partial\gamma}v_c^\gamma\left([\rho_N];\mathbf{r}'\right)\neq\text{constant}$. This proves the main point of this paper:

$$v_{uxc}^\gamma\left(\left[\rho_N\right];\mathbf{r}\right)=v_{uxc}^\gamma\left(\left[\rho_{N-1}^\gamma\right];\mathbf{r}\right)+\text{constant} \tag{34}$$

when ρ_N and ρ_{N-1}^γ are determined by the same external potential $v_{\text{ext}}^\gamma\left(\left[\rho_N\right];\mathbf{r}\right)$.

III. PROOF OF COROLLARY

Let the energy functional $F_N^\gamma[\rho_N]$[12,13] be defined as

$$F_N^\gamma\left[\rho_N\right] = T_N^0\left[\rho_N\right] + \gamma U\left[\rho_N\right] + \gamma E_x\left[\rho_N\right] + E_c^\gamma\left[\rho_N\right]. \tag{35}$$

Then[12,13]

$$\frac{\delta F_N^\gamma\left[\rho_N\right]}{\delta\rho_N\left(\mathbf{r}\right)} + v_{\text{ext}}^\gamma(\left[\rho_N\right];\mathbf{r}) = \mu \tag{36}$$

and

$$\frac{\delta F_N^\gamma\left[\rho_{N-1}^\gamma\right]}{\delta\rho_{N-1}^\gamma\left(\mathbf{r}\right)} + v_{\text{ext}}^\gamma(\left[\rho_N\right];\mathbf{r}) = \mu_{N-1}^\gamma. \tag{37}$$

From the last two equations we find that

$$\frac{\delta F_N^\gamma\left[\rho_N\right]}{\delta\rho_N\left(\mathbf{r}\right)} - \frac{\delta F_N^\gamma\left[\rho_{N-1}^\gamma\right]}{\delta\rho_{N-1}^\gamma\left(\mathbf{r}\right)} = \mu - \mu_{N-1}^\gamma \tag{38}$$

and therefore, using Eqs. (34) and (35), it follows that

$$\frac{\delta T_N^0\left[\rho_N\right]}{\delta\rho_N\left(\mathbf{r}\right)} - \frac{\delta T_N^0\left[\rho_{N-1}^\gamma\right]}{\delta\rho_{N-1}^\gamma\left(\mathbf{r}\right)} = \text{constant} \tag{39}$$

IV. DISCUSSION AND SUMMARY

The relationship in Eq. (34) is valid for $N > 2$. This follows from the step in Eq. (28) where the correlation energy of the $N-1$ electron system is taken as non-zero. In the proof use is made of the N and $N-1$ electron wave functions, hence the proof given here is valid for integer N. The proof for non-integer values of the electrons and will be presented in another paper.

In summary, it was shown that the Hartree-plus-exchange-correlation potential for an integer N-electron ($N > 2$) system differs by a constant form the corresponding potential for an $(N-1)$-electron system if the densities are determined with the same external potential. As a corollary it was shown that the functional derivative of the independent particle kinetic energy functional of the N and $(N-1)$-electron systems also differ by a constant.

REFERENCES

[1] J. Perdew, R. Parr, M. Levy, and J. Balduz, Phys. Rev. Lett. **49**, 1691 (1982).

[2] L. J. Sham and M. Schlüter, Phys. Rev. Lett. **51**, 1888 (1983).

[3] J. Harris and R. O. Jones, J. Phys. F **4**, 1174 (1974).

[4] D. C. Langreth and J. P. Perdew, Solid State Comm. **17**, 1425 (1975).

[5] D. C. Langreth and J. P. Perdew, Phys. Rev. B **15**, 2884 (1977).

[6] O. Gunnarson and B. I. Lundqvist, Phys. Rev. B **13**, 4274 (1976).

[7] P. Hohenberg and W. Kohn, Phys. Rev. B **136**, 864 (1964).

[8] W. Kohn and L. J. Sham, Phys. Rev. A **140**, 1133 (1965).

[9] M. Levy, Natl. Acad. Sci. USA **76**, 6062 (1979).

[10] M. Levy and J. P. Perdew, Phys. Rev. A **32**, 2010 (1985).

[11] A. Görling and M. Levy, Phys. Rev. B **47**, 13105 (1993).

[12] R. G. Parr and W. Yang, *Density Functional Theory of Atoms and Molecules* (Oxford University Press, New York, 1989).

[13] R. M. Dreizler and E. K. U. Gross, *Density Functional Theory* (Springer-Verlag, Berlin, 1990).

[14] R. Jones and O. Gunnarsson, Rew. Mod. Phys. **61**, 689 (1989).

[15] M. Levy and A. Görling, Phys. Rev. A **53**, 3140 (1996).

[16] M. Levy and A. Görling, Phys. Rev. B **53**, 969 (1996).